


**COLUMBIA BORDO  
(Copper and lime based Fungicide)**



**GENERAL:** Bordeaux Fungicide is an old and reliable fungicide that effectively controls various fungus diseases of flowers, ornamentals, trees and fruit.

Copper sulfate and lime based Bordeaux mixture is an outstanding fungicide and bactericide that has been used for decades to control some diseases of tree fruits and nuts, vine fruits, and ornamentals. The ability of Bordeaux mixture to weather the , winter, and spring rains and to adhere to plants makes it an excellent choice for a winter fungicide. this mixture is primarily used as a fungicide to control bacterial leaf spots, blights, anthracnose, downy mildews and cankers. It also acts as a repellent to many Insects and is labeled for use on many vegetable, tree fruit and nut Crops.

Fixed copper fungicide (tribasic copper sulfate, copper oxychloride sulfate, cupric hydroxide) sprays also control some of the same disease-causing organisms as Bordeaux mixture. While fixed copper sprays are much easier to prepare, they are far less persistent and will not withstand winter rains as nicely as Copper sulfate and lime based Bordeaux mixture . Trasco columbia Bodo is most effective and are a better choice to use in spring after the trees have broken dormancy and tender leaves are exposed. It is effective e for certain pathogens such as the leaf curl fungus. Among its many uses are applications in winter to control the overwintering fire blight inoculum in pears and apples, leaf curl and shot hole pathogens in peach and nectarine, downy mildew and powdery mildew fungi in most crops including vegetables and in grapes to control blight and black rot, peacock spot pathogen in olives, walnut blight bacteria in walnut, and black spot fungus in roses. Common Targeted pathogens

Plant	Disease	Application
Grapes	Blight, Black Rot	Apply in Spring and/or early Summer, repeat as needed.
Apples	Blotch, Bitter Rot, Black Rot, Late Scab	Apply in Spring and/or early Summer, repeat as needed
Peaches	Powdery mildews and downy mildew	Apply as prophylactic spray.

Plant	Disease	Application
Iris, Chrysanthemum	Leaf Spot, Blight	Apply as growth starts in Spring. Repeat as needed
Lilies, Dahlias, Tulips	Botrytis Blight	Repeat at 10-day intervals if plant is diseased.
Gladiolus, Hollyhocks, Phlox	Anthracnose, Leaf Spot	Spray after periods of wet weather.
Lilies, Dahlias, Tulips	Botrytis Blight	Repeat at 10-day intervals if plant is diseased.

Gladiolus, Hollyhocks, Phlox	Anthrachnose, Leaf Spot	Spray after periods of wet weather.
Pansy	Downy Mildew	Spray every 2 weeks during growing season.

**Composition:**

Copper 20%

Lime 6%

Inert Carriers 74%

Total 100%

**Dose:** To be used as a foliar Spray @ 3gms/lit of water

**PRECAUTION:**

1. Use the product at the recommended dose only
2. Do not delay the spraying after the tank mix is prepared
3. Spraying should commence either early morning or just before sunset to avoid any scorching
4. The mixture in the tank should be stirred thoroughly before spraying
5. Always store the product in a cool and dry place
6. This product is not compatible with sulphur /lime sulphur

**Manufactured by :Super Agro India Pvt Ltd**

**Address : 168/2, Lenin Sarani, Kolkata-700 013**